

SUPPLEMENTAL MATERIAL

STING-IRF3 Triggers Endothelial Inflammation in Response to Free Fatty Acid-Induced Mitochondrial Damage in Diet-Induced Obesity

Yun Mao, Wei Luo, Lin Zhang, Weiwei Wu, Liangshuai Yuan, Hao Xu, Juhee Song, Keigi Fujiwara, Jun-ichi Abe, Scott A. LeMaire, Xing Li Wang, Ying. H. Shen

From the Shandong University Qilu Hospital Research Center for Cell Therapy, Key Laboratory of Cardiovascular Remodeling and Function Research (Y.M., W.L., W.W., L.Y., H.X., X.L.W.), Qilu Hospital of Shandong University, Jinan 250012, P.R. China; Department of Surgery (Y.M., L.Z., S.A.L., X.L.W., Y.H.S.), Baylor College of Medicine, Houston, TX, USA; Department of Surgery (Y.M., L.Z., S.A.L., X.L.W., Y.H.S.), Texas Heart Institute, Houston, TX, USA; Department of Biostatistics (J.S.), The University of Texas MD Anderson Cancer Center, Houston, TX, USA; and Department of Cardiology - Research, Division of Internal Medicine (K.F., J.-I.A.), The University of Texas MD Anderson Cancer Center, Houston, TX, USA.

Supplemental Figure Legends:

Figure S I. PA induced STING-IRF3 activation and ICAM-1 expression to a similar extent and in a similar pattern in HAECs from different donors. Two batches of HAECs from different donors were treated with PA for 24 hours. **A**, Western blot results indicated that PA induced IRF3 phosphorylation and ICAM-1 expression similarly in the 2 batches of cells. **B-C**, Immunostaining showed that PA induced STING perinuclear translocation (**B**) and IRF3 nuclear translocation (**C**) in both batches of cells.

Figure S II. The STING-IRF3 pathway was activated in the aortas of obese mice.

Wild-type mice were fed either a chow diet (CD) or high-fat diet (HFD) for 12 weeks. **A**, The body weights and free fatty acid (FFA) levels of the 2 groups were compared. **B**, Western blot analysis showed increases in the levels of total IRF3, phospho-IRF3 and ICAM-1 in the aorta of the HFD-fed mice. **C-D**, Immunostaining showed that STING (**C**) and IRF3 (**D**) translocation were increased in the aortic wall of the HFD-fed mice.

Figure S I




Figure S II

